

Rapportage standardisé cavité nasale et sinus paranasaux

Conclusion

Thérapie néoadjuvante : Choisissez une option.

Localisation tumorale : Choisissez une option.

Si autre, spécifiez :

Si plusieurs localisations sinonasales, spécifiez :

Latéralité : Choisissez une option.

Si plusieurs localisations sinonasales, spécifiez :

Focalité : Choisissez une option.

Si multifocale, spécifiez le nombre de foyers :

Si non évaluable, spécifiez :

Type histologique de la tumeur¹ : Choisissez une option.

Si possible, spécifiez le type :

Degré de différenciation : Choisissez une option.

Si non évaluable, spécifiez :

¹ Pour les néoplasies des glandes salivaires accessoires, l'option "Carcinome des glandes salivaires (spécifiez)" doit être sélectionnée. Le sous-type histologique peut ensuite être précisé selon la liste incluse dans le rapportage standardisé "Glandes salivaires principales".

Extension locale : Choisissez une option.

Si une extension locale a été démontrée :

Invasion osseuse/cartilagineuse² : Choisissez une option.

Invasion des tissus mous : Choisissez une option.

Invasion de la base du crâne : Choisissez une option.

Invasion de la peau : Choisissez une option.

Invasion des tissus de l'oeil : Choisissez une option.

Invasion à un autre endroit que ceux décrits ci-dessus : Choisissez une option.

Si invasion à un autre endroit que ceux décrits ci-dessus, spécifiez :

Si l'extension locale ne peut être évaluée, spécifiez :

Invasion lymphovasculaire : Choisissez une option.

S'il est possible de différencier entre invasion des vaisseaux lymphatiques et sanguins, spécifiez : Choisissez une option.

Si non déterminée avec certitude, spécifiez :

Invasion périneurale³ : Choisissez une option.

Diamètre de la plus grande racine nerveuse affectée (*en mm*) : Choisissez une option.

Si non déterminée avec certitude, spécifiez :

² La présence d'une invasion osseuse influence le stade de la tumeur et constitue un facteur pronostic négatif. Une invasion osseuse et/ou cartilagineuse est fréquente pour les carcinomes sinonasaux. L'érosion et la destruction osseuse doivent toutes les deux être rapportées dans le cadre de la classification TNM. Pour plus d'informations, voir 'Note 4 – Extent of invasion' plus bas dans ce document.

³ La présence ou l'absence d'invasion périneurale péri-tumorale peut avoir une influence sur le traitement et le pronostic. Il s'agit d'une invasion périneurale péri-tumorale le long ou au-delà des limites de la tumeur, et donc pas d'une extension intratumorale. Pour plus d'informations, voir 'Note 6 – Perineural invasion' plus bas dans ce document.

Marge minimale définitive saine

(y compris les éventuelles résections ultérieures et sections congelées)⁴ : Choisissez une option.

Marge minimale définitive saine (en mm) :

Précisez la marge la plus proche, si possible :

Si non évaluable, spécifiez :

Classification pTNM (8^{ème} édition du TNM de l'UICC, avec errata)⁵ : pT

Choisissez la classification d'application :

Sinus maxillaire : Choisissez une option.

Cavité nasale et sinus ethmoïdal : Choisissez une option.

Préfixes :

m (multiples tumeurs primaires) : Choisissez une option.

r (récidive) : Choisissez une option.

y (après thérapie néoadjuvante) : Choisissez une option.

⁴ Pour les tumeurs sinonasales, les marges sont généralement rapportées comme positives ou négatives sans qu'il ne soit possible de déterminer la distance par rapport à la marge de résection. Une marge de résection avec du carcinome intraépithélial doit être considérée comme positive pour du carcinome invasif. Pour plus d'informations, voir 'Note 7 – Margin status' plus bas dans ce document.

⁵ La stadification pathologique est basée sur la 'TNM Classification of Malignant Tumours' (8^{ème} édition, UICC). Une description de chaque catégorie pT est disponible dans la 'Note 10 – Pathological staging' plus bas dans ce document.

Champ d'application

Echantillons de résections et de biopsies de tumeurs des muqueuses des cavités nasales et des sinus paranasaux, y compris les tumeurs au bord de la base du crâne.

Remarques

- Pour des résections relatives à des récurrences, le dataset peut être utilisé de manière pragmatique, même si certaines variables peuvent ne pas être applicables ou évaluables.
- Les tumeurs des os et des tissus mous sont traitées dans un dataset distinct de l'ICCR.
- Les néoplasies neuro-endocrines sont traitées dans un dataset distinct.
- Les dissections cervicales et les excisions ganglionnaires sont traitées dans un dataset distinct. Si applicable, l'utilisation de ce dataset doit être combinée.
- Un dataset distinct doit toujours être rempli pour chaque tumeur différente.

Dataset basé sur

Bishop JA, Agaimy A, Bal M, Franchi A, Gallia GL, Kahn S, Rooper L, Stelow E, Weinreb I, Helliwell T, Thompson LDR (2024). *Carcinomas of the Nasal Cavity and Paranasal Sinuses Histopathology Reporting Guide. 2nd edition*. International Collaboration on Cancer Reporting; Sydney, Australia. ISBN: 978-1-922324-52-8.

Note 1 – Tumour site

The sinonasal tract consists of the nasal cavity and the paranasal sinuses (maxillary, ethmoid, frontal, and sphenoid). The nasal cavity can be further subdivided into the nasal septum, floor, lateral wall, and vestibule. Among sinonasal tract carcinomas, the most common site of tumour origin is the maxillary sinus, followed by the nasal cavity and ethmoid sinus. It is rare for carcinomas to arise from the frontal or sphenoid sinuses, except for neuroendocrine tumours in the sphenoid of pituitary origin.¹²⁻¹⁶

The precise tumour site within the sinonasal tract is important to record. First, different staging schemes are utilised for maxillary sinus carcinomas and those arising in the ethmoid sinus or nasal cavity.^{17,18} Second, there is prognostic importance to the tumour location. For example, carcinomas primary to the nasal cavity have been shown to have an improved prognosis over carcinomas primary to the paranasal sinuses, likely because nasal carcinomas give rise to symptoms (e.g., nasal obstruction or epistaxis) and thus come to clinical attention sooner.^{12,16,19,20} In addition, among maxillary sinus carcinomas, those arising from the anterior-inferior portion have a better prognosis than those arising from the superior-posterior portion, likely because the latter group has easier access to structures such as the orbit or skull base.^{17,18} Finally, certain carcinomas are closely associated with specific sinonasal sub-sites. For example, intestinal-type adenocarcinomas and neuroendocrine carcinomas occur most often in the ethmoid sinuses, while squamous cell carcinoma (SCC) occurs most often in the maxillary sinus.²¹⁻²⁴

It is recognised that many carcinomas affect more than one sinonasal anatomic sub-site. In this case, every affected site should be selected.

Note 2 – Histological tumour type

All sinonasal tumours should be classified based on the most recent edition of the WHO Classification of Head and Neck Tumours, 5th edition, 2024 (Table 1).³ The list of histologic types discussed in the chapter on sinonasal tumours in the 5th edition of the WHO does not include salivary gland type tumours or neuroendocrine tumours because they are described in sections devoted to those topics. Tumours of these types should be reported using the this dataset, with reference to the ICCR Carcinomas of the major salivary glands dataset for guidance on histological typing.²⁵ Neuroendocrine neoplasms, specifically carcinomas (small cell and large cell) develop in this site and are recorded here. Neuroendocrine tumours grade 1 and 2 are vanishingly rare, and thus are not specifically included, but can be entered in 'other'.

The sinonasal tract gives rise to a very large and diverse group of malignant tumours.

Accurate tumour typing is important because specific tumour types are associated with different prognoses and, in some cases, different treatments.

Diagnostic accuracy is also expected to take on additional importance in the future as targeted,

⁶ Informations complémentaires ('notes') tirées et adaptées de Bishop JA, Agaimy A, Bal M, Franchi A, Gallia GL, Kahn S, Rooper L, Stelow E, Weinreb I, Helliwell T, Thompson LDR (2024). Carcinomas of the Nasal Cavity and Paranasal Sinuses Histopathology Reporting Guide. 2nd edition. International Collaboration on Cancer Reporting; Sydney, Australia. ISBN: 978-1-922324-52-8.

molecular- based therapies become more prominent. While routine histologic examination has historically been the mainstay for diagnosis, an increasingly large number of sinonasal malignancies require ancillary testing to diagnose (see **NOTE 14 – ANCILLARY STUDIES**). Because these ancillary techniques are not universally available, a diagnosis of ‘Carcinoma, not otherwise specified’ with an explanatory comment may be given if a diagnosis cannot be further refined with the testing methods available to the pathologist.

Table 1: World Health Organization classification of tumours of the nasal cavity, paranasal sinuses and skull base.³

Descriptor	ICD-O codes^a
Carcinomas	
Keratinising squamous cell carcinoma	8071/3
Other squamous cell carcinoma subtypes: Papillary, verrucous, spindle cell, acantholytic, adenosquamous, carcinoma cuniculatum	
Non-keratinising squamous cell carcinoma	8072/3
NUT carcinoma	8023/3
SWI/SNF complex-deficient sinonasal carcinoma	8044/3
Sinonasal lymphoepithelial carcinoma	8082/3
Sinonasal undifferentiated carcinoma	8020/3
Teratocarcinosarcoma	9081/3
HPV-related multiphenotypic sinonasal carcinoma	8483/3
Adenocarcinoma	
Intestinal-type adenocarcinoma	8144/3
Non-intestinal-type adenocarcinoma	8140/3
Neuroendocrine neoplasms	
Small cell neuroendocrine carcinoma	8041/3
Large cell neuroendocrine carcinoma	8013/3
Carcinoma mixed with small cell neuroendocrine carcinoma ^b	8045/3
Carcinoma mixed with large cell neuroendocrine carcinoma ^b	8013/3

^a These morphology codes are from the International Classification of Diseases for Oncology, third edition, second revision (ICD-O-3.2).²⁶ Behaviour is coded /0 for benign tumours; /1 for unspecified, borderline, or uncertain behaviour; /2 for carcinoma in situ and grade III intraepithelial neoplasia; /3 for malignant tumours, primary site; and /6 for malignant tumours, metastatic site. Behaviour code /6 is not generally used by cancer registries. Subtype labels are indented.

^b This terminology is synonymous with the ICD-O terminology of combined small/large cell neuroendocrine carcinomas.

© World Health Organization/International Agency for Research on Cancer. Reproduced with permission.

Note 3 – Histological tumour grade

The applicability of tumour grading in the sinonasal tract is dependent on the histologic type (see Table 2). Most newly-described entities have no established grading scheme, but rather are known to have inherent biologic behaviour (e.g., NUT carcinoma is very aggressive, while human papillomavirus (HPV)-related multiphenotypic sinonasal carcinoma is relatively indolent).^{27,28}

Still, several grading systems for each tumour type are available, with differing merits, and as such, recording which system has been applied is more clinically meaningful (use 'specify' to state the system used), with the ICCR deferring to the WHO Classification current edition for grading guidance and preference.³

Table 2: Applicable grading schemes for sinonasal tumour types.

Histologic tumour type	Grading scheme
Keratinising squamous cell carcinoma	Well-, moderately-, or poorly-differentiated
Non-keratinising squamous cell carcinoma	Not applicable
NUT carcinoma	Not applicable
SWI/SNF complex-deficient sinonasal carcinoma	Not applicable
Sinonasal lymphoepithelial carcinoma	Not applicable
Sinonasal undifferentiated carcinoma	Not applicable
Neuroendocrine carcinoma	High grade
Human papillomavirus (HPV)-related multiphenotypic sinonasal carcinoma	Not applicable
Intestinal-type sinonasal adenocarcinoma	Emerging data to support well-, moderately-, or poorly-differentiated scheme. ²⁹
Non-intestinal type sinonasal adenocarcinoma	Low grade or high grade
Salivary-type adenocarcinoma	See ICCR Carcinoma of major salivary glands dataset. ²⁵
Teratocarcinosarcoma	Not applicable

Note 4 – Extent of invasion

Bone and/or cartilage invasion is a frequent finding in sinonasal carcinomas. Both bone erosion and destruction have to be reported as part of the definition of the primary tumour in the TNM staging system.^{17,18} Soft tissue infiltration and skull base involvement are incorporated into the staging.

Note 5 – Lymphovascular invasion

Lymphovascular invasion consists in the presence of neoplastic cells within an endothelial-lined space, either lymphatic or venous, and should be distinguished from retraction artefact. Immunohistochemical staining for an endothelial marker may help in this distinction.

Lymphovascular invasion is reported in up to 60% of sinonasal SSCs, but its clinical significance at this anatomic site remains to be determined.³⁰

Cases that are still equivocal after taking additional steps may be reported as 'indeterminate' for lymphovascular invasion, but this designation should be sparingly used and it is useful to provide the reason in a comment in the report.

Note 6 – Perineural invasion

The frequency of perineural invasion in sinonasal carcinomas is lower than other head and neck sites, and varies according to the histologic subtype, being most frequent in adenoid cystic carcinoma, sinonasal undifferentiated carcinoma and SSC.^{30,31} In sinonasal carcinomas, perineural invasion is associated with a high rate of positive margins, with variable results with respect to outcome.^{30,32}

Note 7 – Margin status

As endoscopic procedures are now the predominant sinonasal tumour resection, margins are most often sent as numerous separate, fragmented specimens. Therefore, margins can usually only be reported as positive or negative, with distance to margin being impossible to determine. The significance of positive margins has been historically extrapolated from studies on oral cavity tumours,³³⁻³⁵ but there is increasing evidence to support a worse outcome for sinonasal tumours as well.^{32,36-39}

Surface dysplasia and carcinoma in situ are exceedingly rare in sinonasal carcinomas, but secondary surface spread from an invasive carcinoma can be seen.^{28,40} For the purposes of this dataset, a margin with intraepithelial carcinoma should be regarded as positive for invasive carcinoma.

Note 8 – Precursor lesions

It is well established that sinonasal papillomas (especially the inverted and oncocytic subtypes) may give rise to sinonasal carcinomas, most often SSCs but rarely other types.^{41,42} Surface dysplasia is rare in the sinonasal tract, but it is a characteristic precursor lesion in HPV-related multiphenotypic sinonasal carcinoma.²⁸ It has been suggested that some sinonasal non-salivary adenocarcinomas may arise from respiratory epithelial adenomatoid hamartoma or seromucinous hamartoma, but the precursor role of these lesions is unresolved.

Note 9 – Ancillary studies

While keratinising SSC – the most common sinonasal malignancy – can be diagnosed by routine microscopy, ancillary techniques are becoming increasingly necessary to diagnose many sinonasal tumours (see Table 3). If the specific technique is not performed, then a note to that effect should be entered, along with the most likely candidate category (i.e., if NUT immunohistochemistry is not available, state non-keratinising SCC, and suggest it may be in this category).

While parts of this element are deemed core, consideration should be given to temporarily downgrading these to non-core until resources allow.

Table 3: Core and non-core ancillary techniques for each sinonasal tumour type.

Histologic tumour type	Ancillary techniques
Keratinising squamous cell carcinoma	Not needed in most cases (non-core).
Non-keratinising squamous cell carcinoma	Diffuse expression of squamous markers (e.g., p40, CK5/6) required (core). Negative CD99 is useful to exclude adamantinoma-like Ewing sarcoma, and negative NUT to exclude NUT carcinoma (non-core). Many are human papillomavirus (HPV)-related, but HPV testing is not currently required (non-core).
NUT carcinoma	Demonstration of <i>NUT</i> gene rearrangement or positivity with monoclonal antibody against NUT protein is required (core). ⁴³
SWI/SNF complex-deficient sinonasal carcinoma	Loss of expression of either SMARCB1 or SMARCA4 by immunohistochemistry is required (core). ^{44,45}
Sinonasal lymphoepithelial carcinoma	Usually positive for Epstein-Barr virus (EBV) by in situ hybridization. ⁴⁶ Useful but not required (non-core).
Sinonasal undifferentiated carcinoma	Diagnosis of exclusion, so other similar-appearing entities (e.g., non-keratinising squamous cell carcinoma, NUT carcinoma, SWI/SNF complex-deficient sinonasal carcinomas) must be excluded (core). ⁴⁷
Neuroendocrine carcinoma	Positive staining with at least one specific neuroendocrine marker (synaptophysin, chromogranin, INSM1) and an epithelial marker required (core). Other tumours which express these markers must be excluded, e.g., olfactory neuroblastoma, teratocarcinosarcoma.
HPV-related multiphenotypic sinonasal carcinoma	HPV-specific testing (in situ hybridization or PCR) is required (core). Testing should include type 33 which is most common. ⁴⁸
Intestinal-type sinonasal adenocarcinoma	Immunostaining with CDX2 and CK20 is useful but not required (non-core). ⁴⁹
Non-intestinal type sinonasal adenocarcinoma	Often positive for SOX10, S100, and DOG1, with a subset showing nuclear beta-catenin expression, but not required for diagnosis (non-core). ^{50,51}
Salivary-type adenocarcinoma	See ICCR Carcinoma of major salivary glands dataset. ²⁵
Teratocarcinosarcoma	SMARCA4 is often completely or partially lost and beta-catenin staining is frequently nuclear. Useful in a limited sample, but not required to diagnose (non-core). ⁵²

Note 10 – Pathological staging

By Union for International Cancer Control (UICC)/American Joint Committee on Cancer (AJCC) convention,^{17,18} the designation 'T' refers to a primary tumour that has not been previously treated. The symbol 'p' refers to the pathologic classification of the stage, as opposed to the clinical classification, and is based on gross and microscopic examination. pT entails a resection of the primary tumour adequate to evaluate the highest pT category, pN entails removal of nodes adequate to validate lymph node metastasis, and pM implies microscopic examination of distant lesions. There is no pathologic M0 category as this designation requires clinical evaluation and imaging. Clinical classification (cTNM) is usually carried out by the evaluating clinician before treatment during initial evaluation of the patient or when pathologic classification is not possible.

Pathological staging is usually performed after surgical resection of the primary tumour and depends on documentation of the anatomic extent of disease, whether or not the primary tumour has been completely removed. If a biopsied tumour is not resected for any reason (e.g., when technically unfeasible) and if the highest T and N categories or the M1 category of the tumour can be confirmed microscopically, the criteria for pathologic classification and staging have been satisfied even though total removal of the primary cancer was not performed.

Primary tumour (pT)

- TX Primary tumour cannot be assessed
- T0 No evidence of primary tumour
- Tis Carcinoma in situ

Maxillary sinus

- T1 Tumour limited to the mucosa with no erosion or destruction of bone
- T2 Tumour causing bone erosion or destruction, including extension into the hard palate and/or middle nasal meatus, except extension to posterior wall of maxillary sinus and pterygoid plates
- T3 Tumour invades any of the following: bone of posterior wall of maxillary sinus, subcutaneous tissues, floor or medial wall of orbit, pterygoid fossa, or ethmoid sinuses
- T4a Tumour invades any of the following: anterior orbital contents, skin of cheek, pterygoid plates, infratemporal fossa, cribriform plate, sphenoid or frontal sinuses
- T4b Tumour invades any of the following: orbital apex, dura, brain, middle cranial fossa, cranial nerves other than maxillary division of trigeminal nerve (V2), nasopharynx, or clivus

Nasal cavity and ethmoid sinus

- T1 Tumour restricted to one subsite of nasal cavity or ethmoid sinus, with or without bony invasion
- T2 Tumour involves two subsites in a single site or extends to involve an adjacent site within the nasoethmoidal complex, with or without bony invasion
- T3 Tumour extends to invade the medial wall or floor of the orbit, maxillary sinus, palate, or cribriform plate
- T4a Tumour invades any of the following: anterior orbital contents, skin of nose or cheek, minimal extension to anterior cranial fossa, pterygoid plates, sphenoid or frontal sinuses

T4b Tumour invades any of the following: orbital apex, dura, brain, middle cranial fossa, cranial nerves other than V2, nasopharynx, or clivus

TNM Descriptors

For identification of special cases of TNM or pTNM classifications, the 'm' suffix and 'y' and 'r' prefixes are used. Although they do not affect the stage grouping, they indicate cases needing separate analysis.

The 'm' suffix indicates the presence of multiple primary tumours in a single site and is recorded in parentheses: pT(m)NM.

The 'y' prefix indicates those cases in which classification is performed during or following initial multimodality therapy (i.e., neoadjuvant chemotherapy, radiation therapy, or both chemotherapy and radiation therapy). The cTNM or pTNM category is identified by a 'y' prefix. The ycTNM or ypTNM categorises the extent of tumour actually present at the time of that examination. The 'y' categorisation is not an estimate of tumour prior to multimodality therapy (i.e., before initiation of neoadjuvant therapy).

The 'r' prefix indicates a recurrent tumour when staged after a documented disease-free interval, and is identified by the 'r' prefix: rTNM.

For the pN classification of regional lymph nodes, see ICCR Nodal excisions and neck dissection specimens dataset.²

Reporting of pathological staging categories (pT,pN,pM) is based on the evidence available to the pathologist at the time of reporting. As indicated in UICC TNM8 and AJCC TNM8,^{17,18} the final stage grouping of a patient's tumour is based on a combination of pathological staging and other clinical and imaging information.

Pathological staging should not be reported if the submitted specimen is insufficient for definitive staging, especially with biopsy samples (core needle, incisional or excisional). Staging is based on the submitted resection, and even if there is grossly residual disease or there is tumour at the margin, pT staging should only be reported on findings in the resection specimen and/or at operation.^{17,18}

The reference document TNM Supplement: A commentary on uniform use, 5th Edition (C Wittekind et al. editors) may be of assistance when staging.⁵³

Références

- 1 Merlin T, Weston A and Tooher R (2009). Extending an evidence hierarchy to include topics other than treatment: revising the Australian 'levels of evidence'. *BMC Med Res Methodol* 9:34.
- 2 International Collaboration on Cancer Reporting (2024). *Nodal Excisions and Neck Dissection Specimens for Head & Neck Tumours Histopathology Reporting Guide. 2nd edition*. Available from: <https://www.iccr-cancer.org/datasets/published-datasets/head-neck/nodal-excisions/> (Accessed 31st July 2024).
- 3 WHO Classification of Tumours Editorial Board (2024). *Head and Neck Tumours, WHO Classification of Tumours, 5th Edition, Volume 10*. IARC Press, Lyon.
- 4 Nibu K, Sugawara M, Asai M, Ichimura K, Mochiki M, Terahara A, Kawahara N and Asato H (2002). Results of multimodality therapy for squamous cell carcinoma of maxillary sinus. *Cancer* 94(5):1476-1482.
- 5 Samant S, Robbins KT, Vang M, Wan J and Robertson J (2004). Intra-arterial cisplatin and concomitant radiation therapy followed by surgery for advanced paranasal sinus cancer. *Arch Otolaryngol Head Neck Surg* 130(8):948-955.
- 6 Madison Michael L, 2nd, Sorenson JM, Samant S and Robertson JH (2005). The treatment of advanced sinonasal malignancies with pre-operative intra-arterial cisplatin and concurrent radiation. *J Neurooncol* 72(1):67-75.
- 7 Licita L, Suardi S, Bossi P, Locati LD, Mariani L, Quattrone P, Lo Vullo S, Oggionni M, Olmi P, Cantu G, Pierotti MA and Pilotti S (2004). Prediction of TP53 status for primary cisplatin, fluorouracil, and leucovorin chemotherapy in ethmoid sinus intestinal-type adenocarcinoma. *J Clin Oncol* 22(24):4901-4906.
- 8 Meccariello G, Deganello A, Choussy O, Gallo O, Vitali D, De Raucourt D and Georgalas C (2016). Endoscopic nasal versus open approach for the management of sinonasal adenocarcinoma: A pooled-analysis of 1826 patients. *Head Neck* 38 Suppl 1:E2267-2274.
- 9 Roxbury CR, Ishii M, Richmon JD, Blitz AM, Reh DD and Gallia GL (2016). Endonasal Endoscopic Surgery in the Management of Sinonasal and Anterior Skull Base Malignancies. *Head Neck Pathol* 10(1):13-22.
- 10 Llorente JL, Lopez F, Suarez C and Hermesen MA (2014). Sinonasal carcinoma: clinical, pathological, genetic and therapeutic advances. *Nat Rev Clin Oncol* 11(8):460-472.
- 11 Slootweg PJ (2005). Complex head and neck specimens and neck dissections. How to handle them. *J Clin Pathol* 58(3):243-248.
- 12 Ansa B, Goodman M, Ward K, Kono SA, Owonikoko TK, Higgins K, Beitler JJ, Grist W, Wadsworth T, El-Deiry M, Chen AY, Khuri FR, Shin DM and Saba NF (2013). Paranasal sinus squamous cell carcinoma incidence and survival based on Surveillance, Epidemiology, and End Results data, 1973 to 2009. *Cancer* 119(14):2602-2610.
- 13 Robin PE, Powell DJ and Stansbie JM (1979). Carcinoma of the nasal cavity and paranasal sinuses:

incidence and presentation of different histological types. *Clin Otolaryngol Allied Sci* 4(6):431-456.

- 14 Sanghvi S, Khan MN, Patel NR, Yeldandi S, Baredes S and Eloy JA (2014). Epidemiology of sinonasal squamous cell carcinoma: a comprehensive analysis of 4994 patients. *Laryngoscope* 124(1):76-83.
- 15 Takahashi Y, Bell D, Agarwal G, Roberts D, Xie TX, El-Naggar A, Myers JN and Hanna EY (2014). Comprehensive assessment of prognostic markers for sinonasal squamous cell carcinoma. *Head Neck* 36(8):1094-1102.
- 16 Turner JH and Reh DD (2012). Incidence and survival in patients with sinonasal cancer: a historical analysis of population-based data. *Head Neck* 34(6):877-885.
- 17 Brierley JD, Gospodarowicz MK and Wittekind C (eds) (2016). *Union for International Cancer Control. TNM Classification of Malignant Tumours, 8th Edition*, Wiley, USA.
- 18 Amin MB, Edge SB, Greene FL, Byrd DR, Brookland RK, Washington MK, Gershenwald JE, Compton CC, Hess KR, Sullivan DC, Jessup JM, Brierley JD, Gaspar LE, Schilsky RL, Balch CM, Winchester DP, Asare EA, Madera M, Gress DM and Meyer LR (eds) (2017). *AJCC Cancer Staging Manual. 8th ed.*, Springer, New York.
- 19 Dulguerov P, Jacobsen MS, Allal AS, Lehmann W and Calcaterra T (2001). Nasal and paranasal sinus carcinoma: are we making progress? A series of 220 patients and a systematic review. *Cancer* 92(12):3012-3029.
- 20 Thorup C, Sebbesen L, Dano H, Leetmaa M, Andersen M, Buchwald C, Kristensen CA, Bentzen J, Godballe C, Johansen J and Grau C (2010). Carcinoma of the nasal cavity and paranasal sinuses in Denmark 1995-2004. *Acta Oncol* 49(3):389-394.
- 21 Klintenberg C, Olofsson J, Hellquist H and Sokjer H (1984). Adenocarcinoma of the ethmoid sinuses. A review of 28 cases with special reference to wood dust exposure. *Cancer* 54(3):482-488.
- 22 Mitchell EH, Diaz A, Yilmaz T, Roberts D, Levine N, DeMonte F, Hanna EY and Kupferman ME (2012). Multimodality treatment for sinonasal neuroendocrine carcinoma. *Head Neck* 34(10):1372-1376.
- 23 Smith SR, Som P, Fahmy A, Lawson W, Sacks S and Brandwein M (2000). A clinicopathological study of sinonasal neuroendocrine carcinoma and sinonasal undifferentiated carcinoma. *Laryngoscope* 110(10 Pt 1):1617-1622.
- 24 Moreno MA, Roberts DB, Kupferman ME, DeMonte F, El-Naggar AK, Williams M, Rosenthal DS and Hanna EY (2010). Mucosal melanoma of the nose and paranasal sinuses, a contemporary experience from the M. D. Anderson Cancer Center. *Cancer* 116(9):2215-2223.
- 25 International Collaboration on Cancer Reporting (2024). *Carcinomas of the major salivary glands Histopathology Reporting Guide. 2nd edition*. Available from: <https://www.iccr-cancer.org/datasets/published-datasets/head-neck/salivary-glands/> (Accessed 31st July 2024).
- 26 Fritz A, Percy C, Jack A, Shanmugaratnam K, Sobin L, Parkin DM and Whelan S (eds) (2020). *International Classification of Diseases for Oncology, Third edition, Second revision ICD-O-3.2*. Available from: http://www.iacr.com.fr/index.php?option=com_content&view=category&layout=blog&id=100&Itemid=577 (Accessed 16th March 2024).

- 27 Chau NG, Ma C, Danga K, Al-Sayegh H, Nardi V, Barrette R, Lathan CS, DuBois SG, Haddad RI, Shapiro GI, Sallan SE, Dhar A, Nelson JJ and French CA (2020). An Anatomical Site and Genetic-Based Prognostic Model for Patients With Nuclear Protein in Testis (NUT) Midline Carcinoma: Analysis of 124 Patients. *JNCI Cancer Spectr* 4(2):pkz094.
- 28 Bishop JA, Andreassen S, Hang JF, Bullock MJ, Chen TY, Franchi A, Garcia JJ, Gnepp DR, Gomez-Fernandez CR, Ihrler S, Kuo YJ, Lewis JS, Jr., Magliocca KR, Pambuccian S, Sandison A, Uro-Coste E, Stelow E, Kiss K and Westra WH (2017). HPV-related Multiphenotypic Sinonasal Carcinoma: An Expanded Series of 49 Cases of the Tumor Formerly Known as HPV-related Carcinoma With Adenoid Cystic Carcinoma-like Features. *Am J Surg Pathol* 41(12):1690-1701.
- 29 de Gabory L, Waubant A, Verillaud B, Michel J, Malard O, Rumeau C, Jankowski R, Moya-Plana A, Vergez S, Favier V, Mortuaire G, Righini C, Patron V, Thariat J, Dupin C, Coelho J and Bénard A (2023). Multicenter study to assess surgical treatments of 452 sinonasal intestinal-type adenocarcinomas: A REFCOR study. *Eur J Surg Oncol* 49(1):39-46.
- 30 Gil Z, Carlson DL, Gupta A, Lee N, Hoppe B, Shah JP and Kraus DH (2009). Patterns and incidence of neural invasion in patients with cancers of the paranasal sinuses. *Arch Otolaryngol Head Neck Surg* 135(2):173-179.
- 31 Hanna E, Vural E, Prokopakis E, Carrau R, Snyderman C and Weissman J (2007). The sensitivity and specificity of high-resolution imaging in evaluating perineural spread of adenoid cystic carcinoma to the skull base. *Arch Otolaryngol Head Neck Surg* 133(6):541-545.
- 32 Sacks PL, Alvarado R, Sacks R, Kalish L, Campbell R and Harvey R (2022). Prognostic factors and outcomes in minimal access resections of skull base and sinonasal epithelial malignancy. *ANZ J Surg* 92(12):3253-3258.
- 33 Hinni ML, Ferlito A, Brandwein-Gensler MS, Takes RP, Silver CE, Westra WH, Seethala RR, Rodrigo JP, Corry J, Bradford CR, Hunt JL, Stojan P, Devaney KO, Gnepp DR, Hartl DM, Kowalski LP, Rinaldo A and Barnes L (2013). Surgical margins in head and neck cancer: a contemporary review. *Head Neck* 35(9):1362-1370.
- 34 Ch'ng S, Corbett-Burns S, Stanton N, Gao K, Shannon K, Clifford A, Gupta R and Clark JR (2013). Close margin alone does not warrant postoperative adjuvant radiotherapy in oral squamous cell carcinoma. *Cancer* 119(13):2427-2437.
- 35 Sutton DN, Brown JS, Rogers SN, Vaughan ED and Woolgar JA (2003). The prognostic implications of the surgical margin in oral squamous cell carcinoma. *Int J Oral Maxillofac Surg* 32(1):30-34.
- 36 Ferrari M, Mattavelli D, Tomasoni M, Raffetti E, Bossi P, Schreiber A, Orlandi E, Taboni S, Rampinelli V, Gualtieri T, Turri-Zanoni M, Battaglia P, Arosio AD, Bignami M, Tartaro T, Molteni M, Bertazzoni G, Fiaux-Camous D, Jourdain C, Verillaud B, Eu D, Nair D, Moiyadi A, Shetty P, Ghosh-Laskar S, Budrukkar A, Magrini SM, Guillermin S, Faivre S, Piazza C, Gilbert RW, Irish JC, de Almeida JR, Pai P, Herman P, Castelnovo P and Nicolai P (2022). The MUSES*: a prognostic study on 1360 patients with sinonasal cancer undergoing endoscopic surgery-based treatment: *MULTi-institutional collaborative Study on Endoscopically treated Sinonasal cancers. *Eur J Cancer* 171:161-182.
- 37 Arosio AD, Bernasconi DP, Valsecchi MG, Pacifico C, Battaglia P, Bignami M, Ferrari M, Mattavelli D, Rampinelli V, Tomasoni M, Schreiber A, Gualtieri T, Piazza C, Magrini SM, Tartaro T, Molteni M, Lambertoni A, Sileo G, Bossi P, Orlandi E, Bertazzoni G, Fiaux-Camous D, Jourdain C, Verillaud B,

Herman P, Nicolai P, Castelnuovo P and Turri-Zanoni M (2022). Patterns of recurrences in sinonasal cancers undergoing an endoscopic surgery-based treatment: Results of the MUSES* on 940 patients:*MULTi-institutional collaborative Study on Endoscopically treated Sinonasal cancers. *Oral Onco*134:106123.

- 38 Torabi SJ, Spock T, Cardoso B, Chao J, Morse E, Manes RP and Judson BL (2020). Margins in Sinonasal Squamous Cell Carcinoma: Predictors, Outcomes, and the Endoscopic Approach. *Laryngoscope* 130(6):E388-e396.
- 39 Jafari A, Shen SA, Qualliotine JR, Orosco RK, Califano JA and DeConde AS (2019). Impact of margin status on survival after surgery for sinonasal squamous cell carcinoma. *Int Forum Allergy Rhinol* 9(10):1205-1211.
- 40 Shah AA, Jain D, Ababneh E, Agaimy A, Hoschar AP, Griffith CC, Magliocca KR, Wenig BM, Rooper LM and Bishop JA (2020). SMARCB1 (INI-1)-Deficient Adenocarcinoma of the Sinonasal Tract: A Potentially Under-Recognized form of Sinonasal Adenocarcinoma with Occasional Yolk Sac Tumor-Like Features. *Head Neck Pathol* 14(2):465-472.
- 41 Nudell J, Chiosea S and Thompson LD (2014). Carcinoma ex-Schneiderian papilloma (malignant transformation): a clinicopathologic and immunophenotypic study of 20 cases combined with a comprehensive review of the literature. *Head Neck Pathol* 8(3):269-286.
- 42 Maisch S, Mueller SK, Traxdorf M, Weyerer V, Stoehr R, Iro H, Hartmann A and Agaimy A (2020). Sinonasal papillomas: A single centre experience on 137 cases with emphasis on malignant transformation and EGFR/KRAS status in "carcinoma ex papilloma". *Ann Diagn Pathol* 46:151504.
- 43 Haack H, Johnson LA, Fry CJ, Crosby K, Polakiewicz RD, Stelow EB, Hong SM, Schwartz BE, Cameron MJ, Rubin MA, Chang MC, Aster JC and French CA (2009). Diagnosis of NUT midline carcinoma using a NUT-specific monoclonal antibody. *Am J Surg Pathol* 33(7):984-991.
- 44 Agaimy A, Hartmann A, Antonescu CR, Chiosea SI, El-Mofty SK, Geddert H, Iro H, Lewis JS, Jr., Märkl B, Mills SE, Riener MO, Robertson T, Sandison A, Semrau S, Simpson RH, Stelow E, Westra WH and Bishop JA (2017). SMARCB1 (INI-1)-deficient Sinonasal Carcinoma: A Series of 39 Cases Expanding the Morphologic and Clinicopathologic Spectrum of a Recently Described Entity. *Am J Surg Pathol* 41(4):458-471.
- 45 Agaimy A, Jain D, Uddin N, Rooper LM and Bishop JA (2020). SMARCA4-deficient Sinonasal Carcinoma: A Series of 10 Cases Expanding the Genetic Spectrum of SWI/SNF-driven Sinonasal Malignancies. *Am J Surg Pathol* 44(5):703-710.
- 46 Alhazzani H, Alabood S, Alhussien A, Alsadah S, Alghulikah A, Asiri S and Alarifi I (2023). Lymphoepithelial Carcinoma Originated from the Sinonasal Cavity: Case Report and Literature Review. *Case Rep Otolaryngol* 2023:4217102
- 47 Agaimy A, Franchi A, Lund VJ, Skálová A, Bishop JA, Triantafyllou A, Andreasen S, Gnepp DR, Hellquist H, Thompson LDR, Rinaldo A and Ferlito A (2020). Sinonasal Undifferentiated Carcinoma (SNUC): From an Entity to Morphologic Pattern and Back Again-A Historical Perspective. *Adv Anat Pathol* 27(2):51-60.
- 48 Bishop JA, Ogawa T, Stelow EB, Moskaluk CA, Koch WM, Pai SI and Westra WH (2013). Human papillomavirus-related carcinoma with adenoid cystic-like features: a peculiar variant of head and

neck cancer restricted to the sinonasal tract. *Am J Surg Pathol* 37(6):836-844.

- 49 Franchi A, Massi D, Palomba A, Biancalani M and Santucci M (2004). CDX-2, cytokeratin 7 and cytokeratin 20 immunohistochemical expression in the differential diagnosis of primary adenocarcinomas of the sinonasal tract. *Virchows Arch* 445(1):63-67.
- 50 Rooper LM, Thompson LDR, Gagan J, Hwang JSG, London NR, Mikula MW, Stevens TM and Bishop JA (2022). Low-grade non-intestinal-type sinonasal adenocarcinoma: a histologically distinctive but molecularly heterogeneous entity. *Mod Pathol* 35(9):1160-1167.
- 51 Purgina B, Bastaki JM, Duvvuri U and Seethala RR (2015). A Subset of Sinonasal Non-Intestinal Type Adenocarcinomas are Truly Seromucinous Adenocarcinomas: A Morphologic and Immunophenotypic Assessment and Description of a Novel Pitfall. *Head Neck Pathol* 9(4):436-446.
- 52 Rooper LM, Agaimy A, Gagan J, Simpson RHW, Thompson LDR, Trzcinska AM, Ud Din N and Bishop JA (2023). Comprehensive Molecular Profiling of Sinonasal Teratocarcinosarcoma Highlights Recurrent SMARCA4 Inactivation and CTNNB1 Mutations. *Am J Surg Pathol* 47(2):224-233.
- 53 Wittekind C, Brierley JD, van Eycken AL and van Eycken E (eds) (2019). TNM Supplement: A Commentary on Uniform Use, 5th Edition Wiley, USA.